

JOURNAL OF APPROXIMATION THEORY **40**, 173–179 (1984)

One-Codimensional Tchebycheff Subspaces

EITAN LAPIDOT

44A Eder St., Haifa, Israel

Communicated by Oved Shisha

Received August 12, 1980; revised November 29, 1982

A necessary and sufficient condition for a finite dimensional Tchebycheff space of real functions on a real set containing its endpoints to contain a 1-codimensional Tchebycheff subspace is given. Examples of n -dimensional Tchebycheff spaces on closed intervals that do not contain $(n - 1)$ -dimensional Tchebycheff subspaces are given for all $n \geq 3$.

1. INTRODUCTION

By a well-known theorem of Krein, every Tchebycheff space (T -space) of real-valued functions defined on an open interval contains a 1-codimensional T -subspace. In [2] we considered T -spaces on real sets containing at most one endpoint, and gave a necessary and sufficient condition for these T -spaces to contain 1-codimensional T -subspaces.

In this paper we discuss this property for T -spaces on real sets containing both endpoints.

DEFINITION 1 ([1]). The set of the real-valued functions f_1, f_2, \dots, f_k , defined on a real set M , is called a Tchebycheff system (T -system) if

$$U \begin{pmatrix} f_1, f_2, \dots, f_k \\ t_1, t_2, \dots, t_k \end{pmatrix} = \det(f_i(t_j))_{i,j=1}^k \quad (1)$$

has a constant sign for all $t_1, t_2, \dots, t_k \in M$ with $t_1 < t_2 < \dots < t_k$.

If L is the span of a T -system, then it is called a T -space. Clearly, every basis of a T -space is a T -system.

An equivalent definition of a T -space L can be given by means of the number of zeros and the number of sign changes of the elements of L . If there exist $t_1, t_2, \dots, t_{r+1} \in M$ such that $\operatorname{sgn} f(t_i) = -\operatorname{sgn} f(t_{i+1}) \neq 0$ for $i = 1, 2, \dots, r$, we say that f has r sign changes on M . If f has r but not $r + 1$ sign changes, we say that f has exactly r sign changes on M and is denoted

by $S^-(f, M)$. By $Z(f, M)$ we denote the number of the distinct zeros of f in M .

DEFINITION 2 ([5], Lemma 1]). Let L be a k -dimensional space of real-valued functions defined on M . Then L is a T -space if for every $f \in L \setminus \{0\}$, $Z(f, M) \leq k - 1$ and $S^-(f, m) \leq k - 1$.

2. 1-CODIMENSIONAL T -SUBSPACES

We consider T -spaces on a real set M containing its infimum and supremum. Since every 2-dimensional T -space on M contains a positive function, one concludes that it contains a 1-codimensional T -subspace. This is not true in general for T -spaces of higher dimension. In Section 3 we show that for every $n \geq 3$ there exists an n -dimensional T -space on a closed interval that does not contain $(n - 1)$ -dimensional T -subspaces.

We first give a necessary and sufficient condition for a T -space on M to contain a 1-codimensional T -subspace. Let f_1, f_2, \dots, f_{n+2} , $n \geq 1$, be defined on the set M , forming a T -system on it, and let L be their span. If $a = \inf M$ and $b = \sup M$, then

$$L_a = \{f \mid f \in L, f(a) = 0\}, \quad (2)$$

$$L_b = \{f \mid f \in L, f(b) = 0\}, \quad (3)$$

and

$$L_{a,b} = L_a \cap L_b \quad (4)$$

are T -spaces on $M_a = M \setminus \{a\}$, $M_b = M \setminus \{b\}$, and $M_{a,b} = M_a \cap M_b$, respectively.

If L contains a 1-codimensional T -subspace, L' , on M , then

$$L'_a = \{f \mid f \in L', f(a) = 0\}, \quad (2')$$

and

$$L'_b = \{f \mid f \in L', f(b) = 0\} \quad (3')$$

will be 1-codimensional T -subspaces on M_a and M_b , respectively. If $n \geq 2$, then

$$L'_{a,b} = L'_a \cap L'_b \quad (4')$$

will be a 1-codimensional T -subspace of $L_{a,b}$ on $M_{a,b}$.

We now show that the converse is also true.

THEOREM 1. *Let M be a real set containing both its infimum and supremum. Let $\{f_i\}_{i=1}^{n+2}$ be a set of $n+2$ real-valued functions defined on M and forming a T -system on it and let L_a , L_b , and $L_{a,b}$ be defined as above. Then L contains a 1-codimensional T -subspace on M iff*

(i) L_a and L_b have 1-codimensional T -subspaces L'_a and L'_b on M_a and M_b , respectively, and

(ii) if $n \geq 2$, then $L'_{a,b} = L'_a \cap L'_b$ has dimension $n-1$.

Remark. If $L'_{a,b}$ is an $(n-1)$ -dimensional space, then it is a T -space on $M_{a,b}$.

Proof. We need only show the "if part." Let $\{g_1, g_2, \dots, g_n, u, v\}$ be a basis of L such that

$$g_i(a) = g_i(b) = 0, \quad i = 1, 2, \dots, n, \quad (5)$$

$$u(a) = (-1)^n, \quad u(b) = 0, \quad (6)$$

and

$$v(a) = 0, \quad v(b) = 1. \quad (7)$$

Clearly, $L_a = \text{span}\{g_1, g_2, \dots, g_n, v\}$, $L_b = \text{span}\{g_1, g_2, \dots, g_n, u\}$ and $L_{a,b} = \text{span}\{g_1, g_2, \dots, g_n\}$.

Assume first that $n \geq 2$. By our hypothesis, we may assume (replacing u and v by $u - \sum_{i=1}^n a_i g_i$ and $v - \sum_{i=1}^n b_i g_i$ if necessary) that

$$L'_a = \text{span}\{g_1, g_2, \dots, g_{n-1}, v\}, \quad (8)$$

and

$$L'_b = \text{span}\{g_1, g_2, \dots, g_{n-1}, u\}. \quad (9)$$

Clearly

$$L'_{a,b} = L'_a \cap L'_b. \quad (10)$$

We may also assume that

$$U \begin{pmatrix} g_1, g_2, \dots, g_n \\ t_1, t_2, \dots, t_n \end{pmatrix} > 0, \quad \text{whenever } a < t_1 < t_2 < \dots < t_n < b, \quad (11)$$

and

$$U \begin{pmatrix} g_1, g_2, \dots, g_{n-1} \\ t_1, t_2, \dots, t_{n-1} \end{pmatrix} > 0, \quad \text{whenever } a < t_1 < t_2 < \dots < t_{n-1} < b. \quad (12)$$

Hence

$$U \begin{pmatrix} g_1, g_2, \dots, g_{n-1}, v \\ t_1, t_2, \dots, t_{n-1}, t_n \end{pmatrix} > 0, \quad \text{whenever } a < t_1 < t_2 < \dots < t_n \leq b, \quad (13)$$

$$U \begin{pmatrix} g_1, g_2, \dots, g_n, v \\ t_1, t_2, \dots, t_n, t_{n+1} \end{pmatrix} > 0, \quad \text{whenever } a < t_1 < t_2 < \dots < t_{n+1} \leq b, \quad (14)$$

$$U \begin{pmatrix} g_1, g_2, \dots, g_{n-1}, u \\ t_1, t_2, \dots, t_{n-1}, t_n \end{pmatrix} < 0, \quad \text{whenever } a \leq t_1 < t_2 < \dots < t_n < b, \quad (15)$$

also

$$U \begin{pmatrix} g_1, g_2, \dots, g_n, u \\ t_1, t_2, \dots, t_n, t_{n+1} \end{pmatrix} > 0, \quad \text{whenever } a \leq t_1 < t_2 < \dots < t_{n+1} < b. \quad (16)$$

We now show that $\{g_1, g_2, \dots, g_{n-1}, u, v\}$ forms a T -system on M . Consider the function $h = Au + Bv + \sum_{i=1}^{n-1} c_i g_i$. If $AB = 0$, then $Z(h, M) \leq n$ and $S^-(h, M) \leq n$. If $AB < 0$, then it follows from (13) and (15) that $Z(h, M_{a,b}) \leq n-1$ and $Z(h, M) \leq n-1$. Also $S^-(h, M_{a,b}) \leq n-1$ and since $h(a) = (-1)^n A$ and $h(b) = B$, one concludes that $S^-(h, M) \leq n-1$. Similarly, in case $AB > 0$, (14) and (16) imply that $Z(h, M) \leq n$ and $S^-(h, M) \leq n$. Hence $\{g_1, g_2, \dots, g_{n-1}, u, v\}$ is a T -system on M . This completes the proof of the theorem for $n \geq 2$.

For $n = 1$, L is a 3-dimensional T -space. $L_{a,b}$ is a 1-dimensional space, i.e., $L_{a,b} = \text{span}\{g\}$, where $g(a) = g(b) = 0$. Also, $L_a = \text{span}\{g, v\}$, $L_b = \text{span}\{g, u\}$, $L'_a = \text{span}\{v\}$, and $L'_b = \text{span}\{u\}$ are T -spaces on M_a and M_b , and as before $L' = \text{span}\{u, v\}$ is a T -space on M , which completes the proof of the theorem.

Theorem 1 requires that M contain at least $n + 2$ points including its endpoints. We now assume that M has a betweenness property, namely, if $x, y \in M$ with $x < y$, then there exists a point $z \in M$ with $x < z < y$.

Let now $w_a(t) = \max\{\max_{1 \leq i \leq n} |g_i(t)|, |v(t)|\}$ for $t \in M_a$ and $w_b(t) = \max\{\max_{1 \leq i \leq n} |g_i(t)|, |u(t)|\}$ for $t \in M_b$. We define

$$y_i(t) = g_i(t)/w_b(t), \quad i = 1, 2, \dots, n, \quad t \in M_b, \quad (17)$$

$$y_{n+1}(t) = u(t)/w_b(t), \quad t \in M_b \quad (18)$$

$$z_i(t) = g_i(t)/w_a(t), \quad i = 1, 2, \dots, n, \quad t \in M_a, \quad (19)$$

and

$$z_{n+1}(t) = v(t)/w_a(t), \quad t \in M_a. \quad (20)$$

Extend (17)–(20) to M by

$$y_i(b) = \lim_{t \rightarrow b'} y_i(t), \quad i = 1, 2, \dots, n+1, \quad (21)$$

$$z_i(a) = \lim_{t \rightarrow a'} z_i(t), \quad i = 1, 2, \dots, n+1, \quad (22)$$

where $a' = \inf M_a$ and $b' = \sup M_b$ (see [2, 3]).

By [2], a necessary and sufficient condition for L_a and L_b to contain a 1-codimensional T -subspace on M_a and M_b , respectively, is that

$$(y_1(a), \dots, y_{n+1}(a)) \quad \text{and} \quad (y_1(b), \dots, y_{n+1}(b)) \quad (23)$$

are not proportional and also

$$(z_1(a), \dots, z_{n+1}(a)) \quad \text{and} \quad (z_1(b), \dots, z_{n+1}(b)) \quad (24)$$

are not proportional.

Following the technique of [2, Theorem 1], one finds that $L'_a \cap L'_b$ is an $(n-1)$ -dimensional space iff

$$(y_1(b), \dots, y_n(b)) = \pm(z_1(a), \dots, z_n(a)) \neq (0, \dots, 0). \quad (25)$$

We have proved

THEOREM 2. *Let $g_1, g_2, \dots, g_n, u, v$ be as in Theorem 1. Let M have the betweenness property and let $y_1, y_2, \dots, y_{n+1}, z_1, z_2, \dots, z_{n+1}$ be defined by (17)–(22). Then $\text{span}\{g_1, g_2, \dots, g_n, u, v\}$ contains a 1-codimensional T -subspace on M iff (23)–(25) hold.*

Notice that in [2], continuity had been assumed but it can be easily seen that the results of [2] hold without any continuity assumptions.

3. T -SPACES THAT HAVE NO 1-CODIMENSIONAL T -SUBSPACES

In [4], Zielke shows that for every $n > 2$, there exist n -dimensional T -spaces, on closed and on half open intervals that have no Markov basis. We now show, using the same and analogous examples, that for every $n > 2$ there exists an n -dimensional T -space on a closed interval that has no 1-codimensional T -subspace.

Case 1 (n odd). Consider first the $(n-1)$ -dimensional space spanned by

$$f_1(t) = t$$

$$f_i(t) = t^{i-2}(t^2 - 1), \quad i = 2, 3, \dots, n-1,$$

defined on $[-1, 1]$.

This is a T -space [4] which does not contain a 1-codimensional T -subspace [2]. Define $p_0 = 1$,

$$p_i(t) = (1-t)f_i(t), \quad i = 1, 2, \dots, n-1, \text{ on } [-1, 1].$$

$L = \text{span}\{p_0, p_1, \dots, p_{n-1}\}$ is a T -space on $[-1, 1]$ (see [4]), and if L contains a 1-codimensional T -subspace, then L'_1 (in the notation of Theorem 1) would contain a 1-codimensional T -subspace on $[1, 1]$, which is impossible since L'_1 is generated by the f_i 's multiplied by a positive function.

Case 2 (n even ($n > 2$)). Similarly, one can show that $L = \text{span}\{p_0, p_1, \dots, p_{n-1}\}$, where $p_0 = 1$,

$$p_1(t) = 1 - t,$$

and

$$p_i(t) = t^{i-2}(1-t)^2(1+t), \quad i = 2, 3, \dots, n-1,$$

is a T -space on $[-1, 1]$ and since $L_1 = \{p \mid p \in L, p(1) = 0\}$ is a T -space (on $[-1, 1]$) that does not contain a 1-codimensional T -subspace [2, Theorem 1], L does not contain a 1-codimensional T -subspace on $[-1, 1]$.

Zielke shows [6, p. 45], that $\text{span}\{f_1, \dots, f_n, g_1, \dots, g_n\}$, where $f_i(t) = \sin(it)$ and $g_i(t) = \cos(it)$, $i = 1, 2, \dots, n$, is a T -space on $[0, \pi]$, for $n \geq 2$. He proves that this T -space has no Markov basis and raises the question of whether it contains a 1-codimensional T -subspace or not. Applying Theorem 2, one concludes that it contains no such T -subspace.

ACKNOWLEDGMENTS

The author wishes to thank the referee for his helpful suggestions.

REFERENCES

1. S. KARLIN AND J. STUDDEN, "Tchebycheff Systems. With Applications in Analysis and Statistics," Interscience, New York, 1966.
2. E. LAPIDOT, On complete Tchebycheff systems, *J. Approx. Theory* **23** (1978), 324-331.
3. E. LAPIDOT, Tchebycheff subspaces of periodic Tchebycheff spaces, *J. Approx. Theory* **29** (1980), 218-222.

4. R. ZIELKE, Tchebyshev systems that cannot be transformed into a Markov system. *Manuscripta Math.* **17** (1975), 67–71.
5. R. ZIELKE, Alternation properties of Tchebycheff systems and the existence of adjoined functions, *J. Approx. Theory* **10** (1974), 172–184.
6. R. ZIELKE, “Discontinuous Čebyšev Systems,” Springer-Verlag, Berlin, 1979.